

Review

Fashioning the Face: Sensorimotor Simulation Contributes to Facial Expression Recognition

Adrienne Wood,^{1,*} Magdalena Rychlowska,² Sebastian Korb,³
and Paula Niedenthal¹

When we observe a facial expression of emotion, we often mimic it. This automatic mimicry reflects underlying sensorimotor simulation that supports accurate emotion recognition. Why this is so is becoming more obvious: emotions are patterns of expressive, behavioral, physiological, and subjective feeling responses. Activation of one component can therefore automatically activate other components. When people simulate a perceived facial expression, they partially activate the corresponding emotional state in themselves, which provides a basis for inferring the underlying emotion of the expresser. We integrate recent evidence in favor of a role for sensorimotor simulation in emotion recognition. We then connect this account to a domain-general understanding of how sensory information from multiple modalities is integrated to generate perceptual predictions in the brain.

When I wish to find out how wise, or how stupid, or how good, or how wicked is any one, or what are his thoughts at the moment, I fashion the expression of my face, as accurately as possible, in accordance with the expression of his, and then wait to see what thoughts or sentiments arise in my mind or heart, as if to match or correspond with the expression. (Edgar Allan Poe, *The Purloined Letter*, pp. 215–216 [1]).

The Perceptual Challenge of Recognizing Emotion Expressions

Most people are face perception experts [2]. Faces, especially those expressing emotion, automatically capture our attention [3], and we extract the emotional meaning of those faces in a matter of a few hundred milliseconds [4], even subconsciously [5]. Expressions of intense emotion, such as a wide-eyed expression of fear or a toothy grin, may have evolved to be highly distinguishable signals, easily recognizable even from a distance [6]. However, the majority of expressions we encounter are not prototypical expressions. They are instead fleeting, subtle, and somewhat idiosyncratic facial gestures [7] shaped by learning and culture (Box 1). Different emotions, attitudes, and intentions can be communicated with the slight changes in eyebrow position, head tilt, onset dynamic, or lip press [7]. However, we infer emotions from faces effortlessly; the facial expressions of others are unique and useful sources of information about the social environment.

Part of the ability to extract information from faces can be attributed to visual expertise [8]. Experts in any domain develop heightened perceptual sensitivity to diagnostic features of object categories, improving their ability to detect and discriminate between category instances [9]. Humans are undoubtedly experts in faces, which can be considered an especially relevant class

Trends

People's recognition and understanding of others' facial expressions is compromised by experimental (e.g., mechanical blocking) and clinical (e.g., facial paralysis and long-term pacifier use) disruptions to sensorimotor processing in the face.

Emotion perception involves automatic activation of pre- and primary-motor and somatosensory cortices, and the inhibition of activity in sensorimotor networks reduces performance on subtle or challenging emotion recognition tasks.

Sensorimotor simulation flexibly supports not only conceptual processing of facial expression but also, through cross-modal influences on visual processing, the building of a complete percept of the expression.

While automatic and presumably non-conscious, sensorimotor simulation of facial expressions is modulated by the perceiver's social context and motivational state.

¹University of Wisconsin – Madison, 1202 West Johnson Street, Madison, WI 53706, USA

²Cardiff University, Tower Building, Park Place, Cardiff CF10 3AT, UK

³International School for Advanced Studies (SISSA), Via Bonomea 265, 34136 Trieste, Italy

*Correspondence: adrienne.wood@wisc.edu (A. Wood).

Box 1. Culture as a Moderator of Emotion Expression and Sensorimotor Simulation

Despite their relative universality, expressions of emotions and associated display rules are subject to considerable cultural variation [113]. Recently, important cultural variation in facial expression has been explained in terms of the diversity of migratory history of a country. In theory, substantial inward migration should create pressures to interact with strangers and clearly communicate one's intentions. A social environment in which frequently interacting people who lack common emotion language and cultural expectations about which emotion is experienced under a given condition, should, over time, foster a reliance on emotional expression for creating smooth interaction and for building trust.

Consistent with this prediction, two recent studies showed that country-level historical heterogeneity – or the number of source countries that have contributed to the present-day population of a given country over the past 500 years [114] – predicts the endorsement of expressivity norms as well as the production of easily recognizable facial expressions (Figure 1). In particular, individuals from historically heterogeneous cultures reported norms that favored the expression of emotion when an emotion was felt, whereas those from historically homogeneous cultures favored the dissimulation of emotional expression [115]. The former also made facial expressions that on average were more easily recognized by individuals from different cultures [116].

The processes recruited by observers to interpret the meaning of emotional expressions may also be determined by the heterogeneity of migratory history of a country. Indeed, the exchange of eye contact, one possible trigger of sensorimotor simulation [36], varies across cultures and is more frequent in some heterogeneous compared to homogeneous cultures [117]. Because social exchange that is reliant on clear emotion communication would presumably also rely more on sensorimotor simulation, a sensible prediction for future research is that historical heterogeneity determines the degree to which people enact sensorimotor simulation of perceived facial expression, particularly when observing individuals with whom one is not well acquainted.




Figure 1. Country-Level Historical Heterogeneity and Emotional Expressivity. (A) A map depicting the number of source countries that contributed to the present-day population of each nation in the past 500 years [114], with more heterogeneous cultures in green. (B) The cross-cultural variance in emotional expressivity explained by different cultural dimensions ([115] and references therein). Note that historical heterogeneity, or the number of source countries contributing to a country's population, explains 18% of the variance (map generated at <http://ert.co.nz/map/>). Abbreviations: I-C, indicator of collectivism; GLOBE, global leadership and organizational behavior effectiveness survey.

of visual objects; our visual system is honed through experience to be sensitive to subtle differences in relevant facial gestures. In addition to relying on visual expertise, people can infer the emotions felt by another based on context [10]. Emotions tend to occur in specific contexts and in response to particular events. People extract these regularities and develop conceptual emotion knowledge, which may be used to guide **emotion recognition** (see **Glossary**) [11].

In addition to visual and contextual routes to emotion recognition, people might also make use of **sensorimotor simulation**, in which they recreate the motor production of the perceived facial expression in themselves. This subthreshold motor activity in theory triggers partial, often unconscious, activity in other neural systems involved in experiencing the corresponding emotion, and from which the simulator implicitly infers the expresser's internal state. Recent suggestive evidence indicates further that sensorimotor simulation also feeds back to shape the visual percept itself [12].

In the following we build a case for the sensorimotor simulation model of emotion perception. First, we review findings that people automatically simulate perceived facial expressions, and

Glossary

Efference copies: when brain motor cortices execute a motor command, they also generate efference copies of the outgoing motor information to 'alert' sensory (e.g., somatosensory, auditory, etc.) cortices that sensory feedback is about to come in. This alters and may even suppress activity in sensory cortices, which will then respond less to incoming sensory feedback. For example, somatosensory cortices show reduced activity when tactile stimulation is a direct result of self-generated movement compared to externally generated movement [112].

Electromyography (EMG): a technique for measuring muscle activity by recording the electrical signals causing muscle fibers to contract. Facial EMG is the most common technique for quantifying facial mimicry because it can detect even nonvisible muscle contractions that are thought to reflect underlying sensorimotor simulation.

Emotion recognition: this refers to the process by which a person translates the percept of another's facial expression (or other nonverbal signal) into meaningful explicit or implicit knowledge about their underlying emotional, motivational, and/or intentional state.

Facial mimicry: the low-level spontaneous motor reaction commonly observed in perceivers of facial expressions. Typically, the perceiver contracts the same facial muscles involved in the perceived facial expression, but prior knowledge or expectations may influence what expression they 'mimic' [103].

Sensorimotor simulation: processing another's facial expression (or, more generally, any action) often leads to the subthreshold recreation of the motor and somatosensory neural processes involved in producing the facial expression. This simulation may then trigger activation of the associated emotion state, allowing the perceiver to quickly and accurately infer what the expresser is feeling. We have chosen to use the term sensorimotor simulation to emphasize the neural mechanisms of the proposed process. The term is closely related to the concepts of 'embodied simulation' and 'facial mimicry' we avoid the former because it is less

that this simulation contributes to their ability to infer emotional states. We combine these observations with evidence that activating one component of an emotion state, such as a facial expression, triggers other emotion components. Drawing on current thinking about how predictions are integrated into both action and perception, we then propose that sensorimotor simulation augments visual perception of facial expressions. In the process of constructing a prediction, inputs to different sensory modalities feed back to influence one another; we can thus think about sensorimotor simulation as a further source of information in constructing a percept. We hope this account of how people recruit their own emotion experience to understand and make predictions about the emotions of others fosters more productive integration of emotion perception research and domain-general work on perception and cognition [13].

Emotions as Patterns of Adaptive Behavioral, Affective, and Physiological Responses

To infer a role for sensorimotor simulation in emotion perception, we need first to define 'emotion'. Emotions are biologically programmed and functional reactions to relevant stimuli that coordinate brain (e.g., attention, perception, cognition, motivation) and body (e.g., hormones, autonomic nervous system, sensory organs, muscle movements) systems that prepare appropriate behavioral responses [14]. Simply put, if brains evolved to move an organism through space [15], then emotions evolved to organize and direct that movement into an adaptive response [16].

Emotions involve synchronized changes in facial, vocal, bodily gestures, and sympathetic, parasympathetic, and hormonal activity, as well as in distributed neural circuitries of the brain [14]. They also involve recognizable subjective feeling states, and motivation to take a particular action (see [17] for a recent explanation of feeling states being neural predictions of action tendencies). The degree to which components of an emotion system co-occur during a particular instantiation of that emotion, such as anger, is determined by the range of available behaviors [18], emotion regulation, and constraints of culturally learned behavioral and expressive norms (see Box 1 for a discussion of a cultural dimension that shapes emotion expressivity). Nevertheless, the components of a specific emotion system are interdependent and sufficiently organized ([19], but see [20]) such that activating one component, such as a scowling facial expression, will at least partially reactivate other components. Indeed, studies have demonstrated that altering facial expressions, physiological arousal, cognition, or behavior alters the emotional state as a whole ([21–25]; for a review, see [26]). The inter-relatedness of emotion components is important for understanding sensorimotor simulation, but because our review examines the recognition of facial expression, we focus on the possibility that feedback from a facial expression shapes an entire emotion state.

Facial Mimicry as Sensorimotor Simulation

Observers of a facial expression of emotion automatically recreate it (covertly, partially, or completely) [27,28]. We refer to this as 'sensorimotor simulation' to highlight the fact that the perceptual process involves activity in somatosensory and motor systems largely overlapping with those that support the production of the same facial expression. Simulation sometimes, but not always (Box 2), results in facial muscle activity in the perceiver, called **facial mimicry**. Simulation and mimicry can occur even if the expression is irrelevant to the task at hand [29] or is perceived nonconsciously [5,30]. For example, patients with unilateral damage to the occipital visual cortex, who are phenomenally blind in the contralateral visual field, nevertheless show facial mimicry and emotion-congruent changes in physiological arousal when facial expressions are presented to their lesioned hemisphere.

While extant findings point to the existence of imitative capacities early in development [31] and in other primates [32], learning may increase the correspondence between visual input and sensorimotor activity, at least in intentional mimicry [33]. The tendency to simulate or overtly imitate perceived emotion expressions facilitates collective action in a social group and allows an

specified, and use the latter only to refer to contraction of facial muscles during emotion perception.

Transcranial magnetic stimulation (TMS): a noninvasive and precise way to inhibit or amplify the activity of particular cortical brain regions. A magnetic coil is held over the subject's head and a generator delivers electrical pulses to a target brain area.

Box 2. Is Activation of Facial Muscles a Necessary Step in Sensorimotor Simulation?

Blocking or reducing facial mimicry leads to impaired recognition of facial expressions of emotion (see main text). Such findings are typically taken to mean that emotion recognition relies on neural activation of facial muscles and on feedback to neural systems from the activation of facial muscles. An alternative account posits that sensorimotor simulation contributes to emotion recognition via a sensorimotor 'as-if' loop and that activity in pre- and primary-motor cortices does not inevitably result in measurable facial mimicry. Instead, mimicry could constitute a 'spill over' of sensorimotor simulation. On this view, impairment in emotion recognition after blocking of, or interference with, facial mimicry stems not from the reduction of facial mimicry but from the creation of facial feedback that is incongruent with the efference copies reaching somatosensory areas during the internal simulation of the observed facial expression.

In addition to the fact that measurable facial mimicry does not always mediate emotion recognition [52], there are other reasons to expect facial mimicry to be unnecessary for sensorimotor simulation. TMS disrupts emotion processing more when it is applied to primary motor, compared to somatosensory, cortices, suggesting that generating the motor output may be more important than receiving subsequent facial feedback [79]. A study examining how hand gestures augment spatial reasoning found that restricting the hand movements of participants did not affect their reasoning performance, but asking them to generate conflicting motions with their hands reduced accuracy [118]. The ability to simulate gesture production was therefore more important than overtly producing gestures. Facial mimicry manipulations may also be most effective if they disrupt premotor and motor processes, perhaps by requiring participants to generate incongruent movements with facial muscles, rather than simply keeping their faces still.

Facial mimicry has potential metabolic and social costs – for instance, automatically and visibly mimicking another person's anger expression could be dangerous. It would be an inefficient system that required muscle contraction. Nevertheless, we view emotion systems and sensorimotor simulation as involving dynamic brain and body systems for which activation is a continuum rather than dichotomous (e.g., simulating or not; having an emotion or not). Facial motor neurons are an extension of the brain, and therefore if an instance of sensorimotor simulation is particularly activated, activation is more likely to cross the threshold and produce muscle activity. Contraction of facial muscles, while not necessary for emotion recognition, can still therefore augment sensorimotor simulation and subsequent emotion processing benefits.

organism to vicariously learn which stimuli and behaviors are safe, threatening, acceptable, or desirable [34]. Although we argue that sensorimotor simulation is difficult to suppress completely, the probability and intensity of mimicry are sensitive to the perceiver's current motivation to engage with and understand the expresser [35] (Box 3). In particular, measurable facial mimicry is more likely to occur when the perceiver and expresser achieve eye contact, which signals the relevance of the emotion expression to the perceiver [36].

In the present view, facial mimicry reflects, and sometimes augments, sensorimotor simulation, but without being a necessary step in the emotion recognition process. Change in the intensity of facial mimicry (usually measured with **electromyography**, EMG) is only a rough indicator of the degree to which people internally simulate perceived expressions – the absence of measurable facial mimicry does not necessarily indicate an absence of sensorimotor simulation (Box 2).

Facial Expressions Modulate Emotion Experience

How does activity in the somatosensory and motor neural systems of a perceiver contribute to solving the task of inferring an emotion from a perceived facial expression? Empirical evidence confirms that our own facial expressions feed back to modulate our emotional experiences [26]. For instance, covertly inducing participants to smile while they undergo a painful cold-pressor task reduces their physiological arousal and self-reported negative affect [37]. Producing facial expressions can activate other physical components of affect, such as pupil dilation (an indicator of autonomic arousal) [38]. Smiling reduces the rapid neural response to errors (as measured with event-related potentials, ERP) in healthy participants [39] and participants with minor depression [40].

In a complimentary way, inhibiting facial movement can reduce the intensity of emotional experience. For example, paralyzing the corrugator muscle – which produces the furrowed brow of anger and sadness – with injections of botulinum toxin reduces symptoms of depression

Box 3. Social and Contextual Influences on Facial Mimicry and Sensorimotor Simulation

The automaticity of sensorimotor simulation and its putative role in emotion recognition are sometimes challenged because facial mimicry appears to be sensitive to social context (see [119] for a review). Sensorimotor simulation probably contributes more to emotion processing when the perceived expression is ambiguous, when the context does not clearly predict what the expresser may be feeling, or when the perceiver is motivated to know the emotional state of the expresser. Individual characteristics of the perceiver, such as gender, age, empathic tendency, power status, attachment style, and social anxiety, have been shown to influence facial mimicry (reviewed in [10]). Importantly, the extent to which people enact sensorimotor simulation depends on their relationship with the expresser: in-group members and liked others elicit more facial mimicry than do out-group members and disliked others [119].

Emotion perception relies on limited cognitive resources, such as attention and sensorimotor simulation, and it is therefore unsurprising that people engage in it more with people they want or need to understand. Motivational states may moderate automatic simulation at very early stages in the process [35], and people can further inhibit the process by avoiding eye contact [51]. They may also suppress overt, visible facial mimicry in contexts where mirroring the perceived expression would be inappropriate or incongruent with the emotional state of the perceiver: for instance, people can rapidly react to perceived negative expressions with positive expressions [119]. The flexibility of overt facial mimicry is underscored by evidence that mimicry of anger, but not of happiness or sadness, is reduced in close relationships [120]. Thus, measured facial activity in response to another's facial expression may reflect simulation for the sake of emotion recognition, a rapid emotional reaction, or both. These phenomena are difficult to separate, although they may involve partially different neural systems. Consequently, multiple motivational and contextual moderators of facial mimicry demand models that allow top-down modulation of low-level processes (Figure 1J).


[41,42], emotional responding to negative stimuli [43], and activation of the amygdala during intentional expression imitation [44]. Similarly, in patients with facial paralysis, the degree to which the zygomatic muscle – which raises the corners of the lips to create a smile – is immobilized predicts the severity of their depressive symptoms [45]. In sum, the sensorimotor activity involved in making facial expressions partially generates, or at least alters, the associated emotion state.

Simulation and the Recognition of Facial Expression

We have argued that the sensorimotor processes involved in producing a particular facial expression can trigger other components of the emotion system. This implies that simulating another's facial expression can fully or partially activate the associated emotion system in the brain of the perceiver. If so, this should be the basis from which accurate facial expression recognition is achieved [46] (the proposed model is presented in Figure 1).

Substantial research evidence suggests that sensorimotor simulation contributes to accurate and efficient recognition of the specific emotion [47,48], valence [49], intensity [50], and intentionality [51,52] conveyed by facial expressions. Targeted disruption of particular facial muscle contractions most affects recognition of expressions that involve those muscles [53,54]. In the cortical blindness study mentioned above [5], the guesses of the patients as to which emotion was present in their 'blind' visual field were significantly better than chance, and not significantly worse than their categorization performance for expressions shown to their intact visual field. Because the patients displayed facial mimicry in response to facial expressions they perceived nonconsciously, this study provides suggestive evidence that sensorimotor simulation can aide emotion processing even below awareness.

If the coupling between facial expression perception and simulation is consistently increased or inhibited, then individual differences in emotion perception abilities can emerge. For instance, people with mirror-touch synesthesia report experiencing exaggerated sensorimotor simulation and display heightened activity in sensorimotor brain regions when they observe others being touched ([55], see also [56]). These synesthetes perform better than control participants on facial expression recognition tasks, probably because sensorimotor feedback during emotion perception is amplified [57].




Trends in Cognitive Sciences

Figure 1. Simulation and the Recognition of Facial Expression. (A) The female perceiver observes the fearful face of a male expresser. (B) The percept activates the face region of the sensorimotor cortices, and other motor control areas, which may result in facial mimicry. (C) The somatosensory, motor, and premotor cortical activity generates activity in other regions of the brain involved in fear states [26] resulting in either overt cognitive, behavioral, and physiological changes (D) or simulation of those states. (E) This partial activation of the fear state allows the perceiver to explicitly or implicitly recognize the emotion of the expresser. (F) Recent evidence [12] suggests that sensorimotor simulation recursively modulates the clarity of the visual percept. (G) Simulation and (H) emotional responding to a perceived facial expression do not require conscious awareness [5]. (I) Conceptual emotion knowledge contributes to the inferred emotion state [121], while affiliation with and motivation to understand the expresser (J) modulate the likelihood that sensorimotor simulation and facial mimicry will occur (Box 3). While box and arrow diagrams of this sort seem to imply neural modularity and a specific sequence of events, we emphasize the distributed and recursive nature of the emotion perception process, which iteratively recruits visual, somatosensory, motor, and premotor cortices, as well as, subcortically, parts of the limbic system and brainstem (fMRI image from [122]).

On the other hand, a recent study linked deficits in facial mimicry and emotional competence among young and adolescent boys to pacifier use [58]. Specifically, the longer the boys had used a pacifier, the less spontaneous facial mimicry they displayed and the lower their scores on measures of empathy and emotional intelligence years later (Figure 2). No effect of pacifier use on the emotional competence of girls was observed. This difference may be due to greater vulnerability of boys to disruptions of facial expression processing induced by the repetitive inhibition of facial mimicry. By 12 months of age girls on average are superior to boys in their social referencing and use of emotion expressed by caretakers [59,60]. Furthermore, parents talk to girls about emotions more than they talk to boys about this topic [61]. It is thus possible that the dose–response curves are different for boys and girls. Gender and individual differences in emotion socialization notwithstanding, the studies described above suggest that pacifiers can inhibit sensorimotor simulation during an important period in development [62], decoupling perception of facial expressions and automatic mimicry, which has unsurprising consequences for emotion processing (see also [63]).

Other factors that disrupt somatosensory feedback from, or motor output to, the facial muscles are expected to affect emotion recognition. As people age, their sensitivity to bodily feedback declines and may impact on their ability to detect not only their own emotional states [64] but



Trends in Cognitive Sciences

Figure 2. Prolonged Pacifier Use Affects User and Observer Facial Mimicry and Emotion Processing. (A) A study [58] examined the relationship between the duration of pacifier use by children (mean age, 7 years) during infancy and the frequency with which they spontaneously exhibited visible facial mimicry while viewing dynamic emotion expressions. Longer pacifier use was associated with less facial mimicry in boys, but not in girls. (B) In a separate study, duration of pacifier use could predict emotional intelligence scores for adolescent boys, but not for girls. (C) Adults engage in less facial mimicry when viewing babies with mouths occluded by a white box or a pacifier, compared to control images [63] (original images from [123]).

perhaps also the expressions of others [65]. As a more extreme example of individual differences in emotion perception as a result of sensorimotor disruption, preliminary evidence suggests that unilateral facial paralysis alters the ability of patients to detect dynamic changes in asymmetric facial expressions [66]. Specifically, facial paralysis had a greater effect on the sensitivity of patients to facial expressions that began on the same anatomical side of the expresser's face as the paralysis of the patient (see also [45]). At first glance severing or weakening of motor or somatosensory nerves may not be expected to seriously impact on simulation and emotion recognition because we do not consider that overt facial muscle movement is necessary for sensorimotor simulation (Box 2). However, any disruption of the predictive motor-sensory loop could impair sensorimotor simulation. Furthermore, animal models suggest that prolonged facial paralysis can cause the associated motor and somatosensory cortices to shrink or reorganize [67], becoming unavailable for sensorimotor simulation. Whether chronic facial paralysis disrupts emotion recognition or compensatory perceptual strategies can eventually develop is not fully clear (e.g., [68]).

Sensorimotor simulation may be especially useful for emotion recognition when the perceived expression is subtle or ambiguous [69], or the judgment to be made is particularly challenging [46]. Judging the authenticity of a smile, for instance, is a complex task that relies on detecting subtle differences in the morphological specificities and temporal dynamics of the smile. The intensity of automatic facial mimicry corresponds to the degree of smile muscle contraction in the face of the expresser, suggesting that facial mimicry (and presumably sensorimotor simulation) is sensitive to the particular features of an expression [52]. The amount of measurable facial mimicry predicts ratings of the authenticity of a smile [52], and disrupting mimicry impairs the ability of observers to distinguish between videos of people producing spontaneous and intentional smiles [51]. An interesting and currently unanswered question is whether facial mimicry and simulation, being helpful for emotion recognition, could be trained to improve emotional intelligence. Intentional mimicry may be equally challenging for people as intentional posing of facial expressions, as indicated by the systematic differences in faked and spontaneous expressions (e.g., [70]). The level of top-down muscle control involved in deliberate mimicry may disrupt spontaneous sensorimotor influences, and possibly impair or alter emotion recognition, because spontaneous and intentional facial expressions do not recruit the same networks of brain activity [71]. To summarize the current evidence, automatic sensorimotor simulation plays a functional role not only in the recognition of actions ([72], cf. [73]) but also in the processing of emotion expressions.

Neural Bases of Sensorimotor Simulation

Sufficient insight into how sensorimotor simulation is implemented in the brain is now available. The neural circuitry involved in perceiving a facial expression is distributed across cortical and subcortical areas, and involves recursive feed-forward and feed-back processes ([74,75] for review). In addition to visual cortices, visual recognition of another's facial expression reliably activates motor and somatosensory cortices, evidencing the link between perception and sensorimotor simulation [76]. In particular, somatosensory, motor, and premotor cortices are associated with emotion recognition deficits in research on lesion patients [77] and in research using **transcranial magnetic stimulation** (TMS) [76,78,79].

For example, inhibiting the right primary motor (M1) and somatosensory (S1) cortices of female participants with TMS reduced spontaneous facial mimicry and delayed the perception of changes in facial expressions [79]. This finding suggests that disrupting the ability of observers to recruit sensorimotor networks during emotion perception reduces their sensitivity to facial expression dynamics. The application of TMS to the same locations had no effect on mimicry and perception of facial expressions by male participants, representing an interesting gender difference that deserves further investigation [80]. Other studies implicate the premotor cortex [81] and the pre-supplementary area [82] in emotion detection and recognition.

Many simulationist accounts (e.g., [46]) postulate a linear sequence of events in which coarse visual processing is followed by spontaneous facial motor output, which then generates somatosensory processing of facial feedback (see **Box 2** for a discussion of whether facial muscle contractions are necessary for sensorimotor simulation). It is, however, well established that the motor and somatosensory systems are interconnected, and that without the somatosensory system motor output is 'blind' and highly ineffective [83].

Current models of motor functioning reserve a crucial role for sensorimotor integration in the planning and executing of motor acts [84,85]. Instead of solely relying on the time-consuming readout of sensory feedback, the brain uses **efferece copies** of the outgoing motor commands. These corollary discharges of the motor output constitute a form of internal monitoring of the body's state, and actively suppress expected sensory input. Predicted and obtained somatosensory input are continuously compared and mutually subtracted at multiple levels: from the cortex, over subcortical structures, down to the spinal cord. This supports fast and

adaptive regulation of movement and, for example, the ability to distinguish sensory stimulation caused by self-generated movements (e.g., when hitting somebody) from that occurring due to external stimuli (e.g., when being hit).

One of the appeals of simulationist accounts is that they do not require additional systems to the one underlying an organism's own actions. In fact, shared representations (and overlapping neural activation patterns) appear to exist for the production, observation, and imagination of action, as well as for the experience, observation, and imagination of touch [15,86,87]. Simulation of observed action and touch therefore competes with the person's own motor and somatosensory states to influence motor control, physiological responses, and conscious representations [86]. Of relevance for emotion perception, the brain needs to distinguish between simulation of the facial actions of an expresser and its own actual motor output. In the healthy brain, simulation triggered from observation, leading to vicarious activation of motor and somatosensory systems, may be partially inhibited and/or properly attributed to other individuals, through frontal and parietal cortical areas [88]. It has also been suggested that simulation is inhibited by the body's own sensorimotor feedback, based on the observation that limb loss can magnify vicarious motor and somatosensory activation [86]. Inhibitory mechanisms such as these allow a perceiver to recruit preexisting sensorimotor processes for the purpose of understanding the facial expressions of others without leading to widespread and maladaptive mirror-touch synesthesia [55].

Future research will need to isolate the cortical and subcortical motor areas involved in the production of motor signals leading to facial mimicry, and the somatosensory areas involved in processing and comparing motor efference copies and incoming signals from facial feedback. In part, these questions can be tackled by inhibiting specific cortical areas using methods of TMS (e.g., [79]). In the meantime, we can already argue for crosstalk in neural systems that may explain how sensorimotor activity influences visual perception of facial expressions and actions more generally [89].

Emotion Perception and Cross-Modal Sensory Integration


Perceptual inputs in any given modality (e.g., sight, sound, proprioception) are often degraded, unclear, or incomplete, and the brain therefore generates predictions to complete the initial input, sometimes by integrating information from other sensory modalities [90]. Indeed, sensory inputs from one modality (e.g., hearing a scream) feed back to influence perception in another modality (detecting fear in a face), compensating for perceptual ambiguities [91–94]. This is because particular sounds, smells, sights, and tactile cues reliably co-occur in the environment of an individual [95]. A realistically synchronized input to one sensory modality can affect a percept in another modality [92]. For instance, experimental participants are faster to consciously perceive moving visual objects when the direction of the stimuli's movement is congruent with the participants' vestibular input (which informs them about their position in space and co-occurs with changes in their visual field) [91].

Cross-modal perceptual integration is often observable in laboratory experiments involving ambiguous perceptual tasks [94]. Facial expressions are often fleeting and difficult to discriminate; this, coupled with the fact that in many interactions prolonged looking at the face of a social partner is aversive [96], can make the visual signal challenging to process. Emotion perception should therefore benefit from cross-modal integration which allows the formation of a more robust and unambiguous percept [97]. Consistent with this prediction, perceptual adaptation to angry facial expressions transfers to the perception of angry vocalizations, providing suggestive evidence for cross-modal integration of emotion percepts [98].

Another study used a constrictive gel facemask, which provides the wearer with distorted sensory feedback from the face, to alter the typical somatosensory-motor feedback loop of the

participants [12]. While wearing the facemask, participants completed a facial expression discrimination task that did not rely on their conceptual emotion knowledge (as do tasks that have been used in related research) but only on their ability to detect small differences between facial expressions. It was found that facial expression discrimination accuracy was worse for participants wearing the facemask, compared to control participants. This difference did not extend to accuracy on a discrimination task that used non-face stimuli (Figure 3). Speculatively, participants may build a representation of the target facial expression that is partially grounded in sensorimotor activity, and which extends the otherwise limited working memory capacity of the visual system. They can then refer to this representation during the discrimination stage of the task. The representation generated by the individuals wearing the gel facemask may however be distorted and ultimately reduce accuracy. Irrespective of the exact mechanism, the findings suggest that, in addition to informing the judgments of the perceivers about a facial expression (Figure 1C), sensorimotor simulation also contributes directly to visual perceptual processes (Figure 1F).

Ultimately, emotion recognition is a game of prediction because we cannot directly access another's experience. In addition to facilitating accurate emotion perception, simulation may also generate predictions about the future actions and emotions of others [99–102]. Prior expectations about the emotional state of an expresser [103,104], as well as the ongoing affect of the perceiver [105], may also contribute to this predictive process, coloring the visual percept via sensorimotor activation. In this way, simulation becomes another contributor to the multi-modal



Trends in Cognitive Sciences

Figure 3. A Recent Study [12] Revealed that Disrupting Facial Feedback, and Presumably Sensorimotor Simulation, Reduces Accuracy on a Facial Expression Discrimination Task. Half the participants wore a gel facemask that is thought to disrupt the normal motor-somatosensory feedback loop by providing distorted feedback from facial movements. (A) Participants observed a target image, which disappeared then reappeared next to a highly-similar distractor. They responded by indicating which image matched the original target. (B) In addition to completing the task using facial expressions, they also completed the task with images constructed from a morphed horse and cow, which served as a control task that is presumably unaffected by disrupting sensorimotor simulation. (C) The gel facemask reduced accuracy across face stimuli but (D) did not affect performance significantly for the non-face stimuli. This study is the first to use a nonlinguistic, low-level discrimination task to investigate the role of sensorimotor simulation in emotion perception.

percept of the expressed affective state, constructed by predictively combining sensory information with prior experience [106,107].

Concluding Remarks

We have surveyed the substantial evidence that (i) activation of one component of an emotion state (such as a facial expression) often triggers other emotion components, and (ii) perceiving emotion expressions relies in part on automatic simulation of emotion expression production. Connecting these two lines of evidence, we argued that the often-observed sensorimotor simulation process allows a perceiver to access stored knowledge, grounded in the distributed emotion system, about the emotional and motivational states associated with the facial expression. During emotion perception, sensorimotor simulation may partially or fully reactivate related concepts, affective states, and autonomic and behavioral changes, which recursively shape and are shaped by visual information. This form of multi-modal integration can influence the visual percept [12] and generate predictions, a necessary function of all cognitive processes [106].

When considered in the context of current understanding by psychologists of how conceptual knowledge is distributed in the perceptual networks of the brain [108], and how this knowledge is accessed to generate predictions and cross-modal percepts [90], simulationist accounts of emotion recognition do not require any special neural ‘hardware’. Visual emotion perception is augmented, as is performance in other perceptual tasks, by recruiting input from other modalities and recreating emotion states [109]. The process is highly flexible but automatic [35].

Deeper understanding of the complex interplay of vision processing, sensorimotor simulation, learning, and evolved emotion states will inform interventions for people suffering from central [110] or peripheral [45] motor diseases, as well as for disorders associated with mimicry and/or emotion recognition impairments [111]. While many questions remain unanswered (see Outstanding Questions), research continues to generate a progressively more precise and elaborate understanding of how humans can infer complex mental states from even the slightest facial expression.

Acknowledgments

We thank two anonymous reviewers and Lawrence Barsalou for their helpful feedback. A.W. was supported by National Science Foundation (NSF) Graduate Research Fellowship 2012148522, M.R. by European Science Research Council (ESRC)-ORA Grant 505465, and P.N. by NSF BCS-1251101.

References

- Poe, E.A. (1975) The purloined letter. In *The Complete Tales and Poems of Edgar Allan Poe*, pp. 208–222, Random House
- Haxby, J.V. et al. (2000) The distributed human neural system for face perception. *Trends Cogn. Sci.* 4, 223–233
- Pourtois, G. et al. (2013) Brain mechanisms for emotional influences on perception and attention: what is magic and what is not. *Biol. Psychol.* 92, 492–512
- Bijlstra, G. et al. (2010) The social face of emotion recognition: evaluations versus stereotypes. *J. Exp. Soc. Psychol.* 46, 657–663
- Tamietto, M. et al. (2009) Unseen facial and bodily expressions trigger fast emotional reactions. *Proc. Natl. Acad. Sci. U.S.A.* 106, 17661–17666
- Smith, F.W. and Schyns, P.G. (2009) Smile through your fear and sadness transmitting and identifying facial expression signals over a range of viewing distances. *Psychol. Sci.* 20, 1202–1208
- Jack, R.E. et al. (2014) Dynamic facial expressions of emotion transmit an evolving hierarchy of signals over time. *Curr. Biol.* 24, 187–192
- Pollak, S.D. et al. (2009) Development of perceptual expertise in emotion recognition. *Cognition* 110, 242–247
- Folstein, J.R. et al. (2013) Category learning increases discriminability of relevant object dimensions in visual cortex. *Cereb. Cortex* 23, 814–823
- Seibt, B. et al. (2015) Facial mimicry in its social setting. *Front. Psychol.* 6, 1122
- Marchi, F. and Newen, A. (2015) Cognitive penetrability and emotion recognition in human facial expressions. *Front. Psychol.* 6, 828
- Wood, A. et al. (2015) Altering sensorimotor feedback disrupts visual discrimination of facial expressions. *Psychon. Bull. Rev.* Published online November 5, 2015. <http://dx.doi.org/10.3758/s13423-015-0974-5>
- Barsalou, L.W. (2013) Mirroring as pattern completion inferences within situated conceptualizations. *Cortex* 49, 2951–2953
- Panksepp, J. (2005) Affective consciousness: core emotional feelings in animals and humans. *Conscious. Cogn.* 14, 30–80
- Wolpert, D.M. et al. (2003) A unifying computational framework for motor control and social interaction. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 358, 593–602
- Lench, H.C. et al. (2015) A functionalist manifesto: goal related emotions from an evolutionary perspective. *Emotion Rev.* 7, 90–98

Outstanding Questions

Does the historical heterogeneity of a culture predict the degree to which people employ sensorimotor simulation (Box 1)?

To what degree do primary sensorimotor cortices, pre- and supplementary-motor cortices, basal ganglia, and sub-cortical emotion systems each contribute to recognizing facial expressions?

What role does learning play in the link between sensorimotor simulation and emotion recognition? If simulation is not completely ‘hard-wired’, how do infants learn to map visually perceived facial expressions to their own patterns of muscle activity? Do they become more accurate simulators by correcting errors in their simulation-based predictions of others’ facial expressions?

What accounts for occasionally observed gender differences in facial mimicry, emotion recognition ability, and vulnerability of these processes to interference?

17. Lowe, R. and Ziemke, T. (2011) The feeling of action tendencies: on the emotional regulation of goal-directed behavior. *Front. Psychol.* 2, 346
18. Löw, A. et al. (2015) When threat is near, get out of here: dynamics of defensive behavior during freezing and active avoidance. *Psychol. Sci.* 26, 1706–1716
19. Mauss, I.B. et al. (2005) The tie that binds? Coherence among emotion experience, behavior, and physiology. *Emotion* 5, 175–190
20. Evers, C. et al. (2014) Emotion response coherence: a dual-process perspective. *Biol. Psychol.* 98, 43–49
21. Michalak, J. et al. (2014) Sitting posture makes a difference—embodiment effects on depressive memory bias. *Clin. Psychol. Psychother.* 21, 519–524
22. Harmon-Jones, E. and Peterson, C.K. (2009) Supine body position reduces neural response to anger evocation. *Psychol. Sci.* 20, 1209–1210
23. Gray, M.A. et al. (2011) Emotional appraisal is influenced by cardiac afferent information. *Emotion* 12, 180–191
24. Harrison, N.A. et al. (2010) The embodiment of emotional feelings in the brain. *J. Neurosci.* 30, 12878–12888
25. Hawk, S.T. et al. (2012) Face the noise: embodied responses to nonverbal vocalizations of discrete emotions. *J. Pers. Soc. Psychol.* 102, 796
26. Price, T.F. and Harmon-Jones, E. (2015) Embodied emotion: the influence of manipulated facial and bodily states on emotive responses. *Wiley Interdiscip. Rev. Cogn. Sci.* 6, 461–473
27. Krumbhuber, E.G. et al. (2014) Facial mimicry of spontaneous and deliberate Duchenne and non-Duchenne smiles. *J. Nonverbal Behav.* 38, 1–11
28. Korb, S. et al. (2010) Timing and voluntary suppression of facial mimicry to smiling faces in a Go/NoGo task – an EMG study. *Biol. Psychol.* 85, 347–349
29. Lee, T.W. et al. (2008) Controlling emotional expression: behavioral and neural correlates of nonimitative emotional responses. *Cereb. Cortex* 18, 104–113
30. Bornemann, B. et al. (2012) Can you feel what you do not see? Using internal feedback to detect briefly presented emotional stimuli. *Int. J. Psychophysiol.* 85, 116–124
31. Geangu, E. et al. (2016) Three-year-olds' rapid facial EMG responses to emotional facial expressions and body postures. *J. Exp. Child Psychol.* 144, 1–14
32. Mancini, G. et al. (2013) Rapid facial mimicry in geladas. *Sci. Rep.* 3, 1527
33. Yee, E. et al. (2013) Manual experience shapes object representations. *Psychol. Sci.* 24, 909–919
34. Niedenthal, P.M. and Brauer, M. (2012) Social functionality of human emotion. *Annu. Rev. Psychol.* 63, 259–285
35. Carr, E.W. and Winkielman, P. (2014) When mirroring is both simple and 'smart': how mimicry can be embodied, adaptive, and non-representational. *Front. Hum. Neurosci.* 8, 505
36. Rychlowska, M. et al. (2012) From the eye to the heart: eye contact triggers emotion simulation. In *Proceedings of the 4th Workshop on Eye Gaze in Intelligent Human Machine Interaction, Article 5, Association for Computing Machinery.*
37. Kraft, T.L. and Pressman, S.D. (2012) Grin and bear it: the influence of manipulated facial expression on the stress response. *Psychol. Sci.* 23, 1372–1378
38. Lee, I.S. et al. (2013) An amplification of feedback from facial muscles strengthened sympathetic activations to emotional facial cues. *Auton. Neurosci.* 179, 37–42
39. Wiswede, D. et al. (2009) Embodied emotion modulates neural signature of performance monitoring. *PLoS ONE* 4, 1–6
40. Lin, W. et al. (2015) Is it helpful for individuals with minor depression to keep smiling? An event-related potentials analysis. *Soc. Behav. Pers.* 43, 383–396
41. Finzi, E. and Rosenthal, N.E. (2014) Treatment of depression with onabotulinumtoxin A: a randomized, double-blind, placebo controlled trial. *J. Psychiatr. Res.* 52, 1–6
42. Wollmer, M.A. et al. (2012) Facing depression with botulinum toxin: a randomized controlled trial. *J. Psychiatr. Res.* 46, 574–581
43. Davis, J.I. et al. (2010) The effects of BOTOX injections on emotional experience. *Emotion* 10, 433
44. Hennenlotter, A. et al. (2009) The link between facial feedback and neural activity within central circuitries of emotion – new insights from botulinum toxin-induced denervation of frown muscles. *Cereb. Cortex* 19, 537–542
45. VanSwearingen, J.M. et al. (1999) Specific impairment of smiling increases the severity of depressive symptoms in patients with facial neuromuscular disorders. *Aesthet. Plast. Surg.* 23, 416–423
46. Niedenthal, P.M. et al. (2010) The simulation of smiles (sims) model: embodied simulation and the meaning of facial expression. *Behav. Brain Sci.* 33, 417–433
47. Künecke, J. et al. (2014) Facial EMG responses to emotional expressions are related to emotion perception ability. *PLoS ONE* 9, e84053
48. Ipser, A. and Cook, R. (2015) Inducing a concurrent motor load reduces categorization precision for facial expressions. *J. Exp. Psychol. Hum. Perception Perform.* Published online November 20, 2015. <http://dx.doi.org/10.1037/xhp0000177>
49. Hyniewska, S. and Sato, W. (2015) Facial feedback affects valence judgments of dynamic and static emotional expressions. *Front. Psychol.* 6, 291
50. Lobmaier, J.S. and Fischer, M.H. (2015) Facial feedback affects perceived intensity but not quality of emotional expressions. *Brain Sci.* 5, 357–368
51. Rychlowska, M. et al. (2014) Blocking mimicry makes true and false smiles look the same. *PLoS ONE* 9, e90876
52. Korb, S. et al. (2014) The perception and mimicry of facial movements predict judgments of smile authenticity. *PLoS ONE* 9, e99194
53. Ponari, M. et al. (2012) Mapping correspondence between facial mimicry and emotion recognition in healthy subjects. *Emotion* 12, 1398
54. Oberman, L.M. et al. (2007) Face to face: blocking facial mimicry can selectively impair recognition of emotional expressions. *Soc. Neurosci.* 2, 167–178
55. Blakemore, S.J. et al. (2005) Somatosensory activations during the observation of touch and a case of vision–touch synaesthesia. *Brain* 128, 1571–1583
56. Maister, L. et al. (2013) I feel your fear: shared touch between faces facilitates recognition of fearful facial expressions. *Emotion* 13, 7
57. Banissy, M.J. et al. (2011) Superior facial expression, but not identity recognition, in mirror-touch synesthesia. *J. Neurosci.* 31, 1820–1824
58. Niedenthal, P.M. et al. (2012) Negative relations between pacifier use and emotional competence. *Basic Appl. Soc. Psychol.* 34, 387–394
59. Mumme, D.L. et al. (1996) Infants' responses to facial and vocal emotional signals in a social referencing paradigm. *Child Dev.* 67, 3219–3237
60. Rosen, W.D. et al. (1992) An experimental investigation of infant social referencing: mothers' messages and gender differences. *Dev. Psychol.* 28, 1172–1178
61. Fivush, R. et al. (2000) Gender differences in parent–child emotion narratives. *Sex Roles* 42, 233–253
62. Bruderer, A.G. et al. (2015) Sensorimotor influences on speech perception in infancy. *Proc. Natl. Acad. Sci. U.S.A.* 112, 13531–13536
63. Rychlowska, M. et al. (2014) Pacifiers disrupt adults' responses to infants' emotions. *Basic Appl. Soc. Psychol.* 36, 299–308
64. Mendes, W.B. (2010) Weakened links between mind and body in older age: the case for maturational dualism in the experience of emotion. *Emotion Rev.* 2, 240–244
65. Ebner, N.C. et al. (2011) Age and emotion affect how we look at a face: Visual scan patterns differ for own-age versus other-age emotional faces. *Cogn. Emotion* 25, 983–997
66. Korb, S. et al. (2016) Facial mimicry is asymmetrical in an anatomical way. *JAMA Facial Plast. Surg.* <http://dx.doi.org/10.1001/jamafacial.2015.2347>
67. Urrego, D. et al. (2015) Layer 5 pyramidal neurons' dendritic remodeling and increased microglial density in primary motor

- cortex in a murine model of facial paralysis. *Biomed. Res. Int.* 2015, 482023
68. Bate, S. *et al.* (2013) First report of generalized face processing difficulties in möbius sequence. *PLoS ONE* 8, e62656
 69. Beffara, B. *et al.* (2012) Enhanced embodied response following ambiguous emotional processing. *Cogn. Process.* 13, 103–106
 70. Krumhuber, E.G. and Manstead, A.S.R. (2009) Can Duchenne smiles be feigned? New evidence on felt and false smiles. *Emotion* 9, 807–820
 71. Hopf, H.C. *et al.* (1992) Localization of emotional and volitional facial paresis. *Neurology* 42, 1918–1923
 72. Hamilton, A.F. *et al.* (2006) Where does your own action influence your perception of another person's action in the brain? *NeuroImage* 29, 524–535
 73. Press, C. and Cook, R. (2015) Beyond action-specific simulation: domain-general motor contributions to perception. *Trends Cogn. Sci.* 19, 176–178
 74. Atkinson, A.P. and Adolphs, R. (2011) The neuropsychology of face perception: beyond simple dissociations and functional selectivity. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 366, 1726–1738
 75. Sato, W. *et al.* (2015) Spatiotemporal neural network dynamics for the processing of dynamic facial expressions. *Sci. Rep.* 5, 12432
 76. Pitcher, D. *et al.* (2008) Transcranial magnetic stimulation disrupts the perception and embodiment of facial expressions. *J. Neurosci.* 28, 8929–8933
 77. Adolphs, R. *et al.* (2000) A role for somatosensory cortices in the visual recognition of emotion as revealed by three-dimensional lesion mapping. *J. Neurosci.* 20, 2683–2690
 78. Pourtois, G. *et al.* (2004) Dissociable roles of the human somatosensory and superior temporal cortices for processing social face signals. *Eur. J. Neurosci.* 20, 3507–3515
 79. Korb, S. *et al.* (2015) Gender differences in the neural network of facial mimicry of smiles – an rTMS study. *Cortex* 70, 101–114
 80. Stel, M. and van Knippenberg, A. (2008) The role of facial mimicry in the recognition of affect. *Psychol. Sci.* 19, 984–985
 81. Balconi, M. and Bortolotti, A. (2013) Conscious and unconscious face recognition is improved by high-frequency rTMS on pre-motor cortex. *Conscious. Cogn.* 22, 771–778
 82. Rochas, V. *et al.* (2012) Disrupting pre-SMA activity impairs facial happiness recognition: an event-related TMS study. *Cereb. Cortex* 23, 1517–1525
 83. Cole, J.D. and Sedgwick, E.M. (1992) The perceptions of force and of movement in a man without large myelinated sensory afferents below the neck. *J. Physiol.* 449, 503–515
 84. Flanders, M. (2011) What is the biological basis of sensorimotor integration? *Biol. Cybern.* 104, 1–8
 85. Requarth, T. *et al.* (2014) A role for mixed corollary discharge and proprioceptive signals in predicting the sensory consequences of movements. *J. Neurosci.* 34, 16103–16116
 86. Case, L.K. *et al.* (2015) Common coding and dynamic interactions between observed, imagined, and experienced motor and somatosensory activity. *Neuropsychologia* 79, 233–245
 87. de Vignemont, F. (2013) Shared body representations and the 'whose' system. *Neuropsychologia* 55, 128–136
 88. Brass, M. *et al.* (2005) The inhibition of imitative and overlearned responses: a functional double dissociation. *Neuropsychologia* 43, 89–98
 89. Orgs, G. *et al.* (2016) Constructing visual perception of body movement with the motor cortex. *Cereb. Cortex* 26, 440–449
 90. Driver, J. and Noesselt, T. (2008) Multisensory interplay reveals crossmodal influences on 'sensory-specific' brain regions, neural responses, and judgments. *Neuron* 57, 11–23
 91. Salomon, R. *et al.* (2015) Balancing awareness: vestibular signals modulate visual consciousness in the absence of awareness. *Conscious. Cogn.* 36, 289–297
 92. Hu, B. and Knill, D.C. (2010) Kinesthetic information disambiguates visual motion signals. *Curr. Biol.* 20, R436–R437
 93. Dieter, K.C. *et al.* (2014) Kinesthesia can make an invisible hand visible. *Psychol. Sci.* 25, 66–75
 94. Taylor-Clarke, M. *et al.* (2002) Vision modulates somatosensory cortical processing. *Curr. Biol.* 12, 233–236
 95. Roseboom, W. *et al.* (2013) The cross-modal double flash illusion depends on featural similarity between cross-modal inducers. *Sci. Rep.* 3, 3437
 96. Helminen, T.M. *et al.* (2011) Eye contact and arousal: the effects of stimulus duration. *Biol. Psychol.* 88, 124–130
 97. Jessen, S. and Kotz, S.A. (2013) On the role of crossmodal prediction in audiovisual emotion perception. *Front. Hum. Neurosci.* 7, 368
 98. Pye, A. and Bestelmeyer, P.E.G. (2015) Evidence for a supramodal representation of emotion from cross-modal adaptation. *Cognition* 134, 245–251
 99. Genschow, O. and Brass, M. (2015) The predictive chameleon: evidence for anticipated social action. *J. Exp. Psychol. Hum. Percept. Perform.* 41, 265–268
 100. Kaufman, J. and Johnston, P.J. (2014) Facial motion engages predictive visual mechanisms. *PLoS ONE* 9, e91038
 101. Palumbo, L. and Jellema, T. (2013) Beyond face value: does involuntary emotional anticipation shape the perception of dynamic facial expressions? *PLoS ONE* 8, e56003
 102. Hudson, M. *et al.* (2016) One step ahead: the perceived kinematics of others' actions are biased towards expected goals. *J. Exp. Psychol. Gen.* 145, 1–7
 103. Halberstadt, J.B. and Niedenthal, P.M. (2001) Effects of emotion concepts on perceptual memory for emotional expressions. *J. Pers. Soc. Psychol.* 81, 587–598
 104. Halberstadt, J. *et al.* (2009) Emotional conception how embodied emotion concepts guide perception and facial action. *Psychol. Sci.* 20, 1254–1261
 105. Niedenthal, P.M. *et al.* (2001) When did her smile drop? Facial mimicry and the influences of emotional state on the detection of change in emotional expression. *Cogn. Emotion* 15, 853–864
 106. Clark, A. (2014) Perceiving as predicting. In *Perception and its Modalities* (Stokes, D. *et al.*, eds), pp. 23–43, Oxford University Press
 107. Cattaneo, L. *et al.* (2011) One's motor performance predictably modulates the understanding of others' actions through adaptation of premotor visuo-motor neurons. *Soc. Cogn. Affect. Neurosci.* 6, 301–310
 108. Rogers, T.T. and Cox, C.R. (2015) The neural bases of conceptual knowledge: revisiting a golden-age hypothesis in the era of cognitive neuroscience. In *The Wiley Handbook on The Cognitive Neuroscience of Memory* (Addis, D.R. *et al.*, eds), pp. 60–83, John Wiley & Sons
 109. Winkielman, P. (2015) The coherent and fluent mind: how unified consciousness is constructed from cross-modal inputs via integrated processing experiences. *Front. Psychol.* 6, 83
 110. Mermillod, M. *et al.* (2011) Embodying emotional disorders: new hypotheses about possible emotional consequences of motor disorders in Parkinson's disease and Tourette's syndrome. *ISRN Neurol.* 2011, 306918
 111. Neufeld, J. *et al.* (2015) Spontaneous facial mimicry is modulated by joint attention and autistic traits. *Autism Res.* Published online October 7, 2015. <http://dx.doi.org/10.1002/aur.1573>
 112. Shergill, S.S. *et al.* (2013) Modulation of somatosensory processing by action. *NeuroImage* 70, 356–362
 113. Jack, R.E. *et al.* (2012) Facial expressions of emotion are not culturally universal. *Proc. Natl. Acad. Sci. U.S.A.* 109, 7241–7244
 114. Putterman, L. and Weil, D.N. (2010) Post-1500 population flows and the long run determinants of economic growth and inequality. *Q. J. Econ.* 125, 1627–1682
 115. Rychlowska, M. *et al.* (2015) Heterogeneity of long-history migration explains cultural differences in reports of emotional expressivity and the functions of smiles. *Proc. Natl. Acad. Sci. U.S.A.* 112, E2429–E2436
 116. Wood, A. *et al.* (2016) Heterogeneity of long-history migration predicts emotion recognition accuracy. *Emotion* <http://dx.doi.org/10.1037/emo0000137>

117. Akechi, H. *et al.* (2013) Attention to eye contact in the West and East: autonomic responses and evaluative ratings. *PLoS ONE* 8, e59312
118. Hegarty, M. *et al.* (2005) The role of gestures in mental animation. *Spat. Cogn. Comput.* 5, 333–356
119. Hess, U. and Fischer, A. (2014) Emotional mimicry: why and when we mimic emotions. *Soc. Pers. Psychol. Compass* 8, 45–57
120. Häfner, M. and Ijzerman, H. (2011) The face of love spontaneous accommodation as social emotion regulation. *Pers. Soc. Psychol. Bull.* 37, 1551–1563
121. Hess, U. and Harel, S. (2015) The influence of context on emotion recognition in humans. In *2015 11th IEEE International Conference and Workshops on Automatic Face and Gesture Recognition (FG 2015)*, pp. 705–710, IEEE
122. Fullana, M.A. *et al.* (2015) Neural signatures of human fear conditioning: an updated and extended meta-analysis of fMRI studies. *Mol. Psychiatry* Published online June 30, 2015. <http://dx.doi.org/10.1038/mp.2015.88>
123. Gil, S. *et al.* (2011) The evaluation of emotional facial expressions in early postpartum depression mood: a difference between adult and baby faces? *Psychiatry Res.* 186, 281–286